

Preventive Measures in Cardiology: Behavioral and Environmental Determinants of Cardiovascular Disease

Hung Quoc To
Toquochung2010@yahoo.com

ABSTRACT

Cardiovascular disease (CVD) remains the leading cause of global mortality. As of 2017, more than 17 million deaths were attributed to CVD, with over 360 million disability-adjusted life years lost, making CVD the highest-burden disease worldwide. As this burden continues to rise, it is essential to understand how behavioral and environmental conditions predispose individuals to CVD within socioeconomic and environmental contexts. In this review, the term “lifestyle factors” refers to health behaviors influenced by socioeconomic and environmental conditions. Using a health equity lens, this review examines how behavioral and environmental factors are linked to structural disparities, contributing to unequal distributions of cardiovascular risk.

Key risk factors of CVD include hypertension, diabetes, obesity, poor diet, physical inactivity, tobacco use, and alcohol use. Prevention strategies focus on health behavior modifications that may address multiple risk factors simultaneously. Sustained physical activity (PA) has been shown to lower cardiovascular mortality. Likewise, consumption of whole grains, fruits, vegetables, proteins, and omega-3 fatty acids may decrease cardiovascular mortality. However, the ability to adopt protective behaviors and avoid detrimental exposures varies across populations due to distinct neighborhood conditions, environmental conditions, and resource availability. Therefore, cardiovascular risk is often observed in structurally disadvantaged settings with increased exposure and limited access to protective conditions.

CVD is a multifactorial disease characterized by interactions among behavioral, biological, and environmental factors. Prevention strategies that integrate individual behaviors with social and environmental conditions may substantially alleviate the global burden of CVD and reduce cardiovascular inequities.

Keywords: Cardiovascular disease (CVD), population health, behavioral and environmental determinants, chronic exposure, health equity, physiological stress accumulation

INTRODUCTION

Chest pain, fatigue, and shortness of breath are symptoms of cardiovascular disease. Despite medical advances, the global burden of CVD continues to escalate, with 20.5 million deaths recorded in 2021.^{1,2,3} Heart disease refers to multiple conditions, such as coronary artery disease (CAD), cardiomyopathy, heart failure (HF), and rheumatic heart disease.⁴ These conditions occur alongside an unequal distribution of modifiable risk factors. Behavioral determinants such as PA, diet, sleep duration, and tobacco use do not occur independently. They are influenced by and constrained within both socioeconomic and environmental contexts. These factors vary across social position, including race, socioeconomic status, gender, age, and housing stability. They influence an individual's pollution exposure, access to safe activity spaces, cumulative stress burden, and preventive care, which in turn correlate with cardiovascular outcomes.⁵ CAD, characterized by the accumulation of plaque in the coronary arteries, illustrates how these conditions may culminate in adverse outcomes.⁶ CAD mortality is associated with a lack of physical activity, with physically inactive individuals experiencing significantly higher risks of premature CVD and hypertension.⁷ This pattern differs based on neighborhood safety, access to recreational spaces, as well as transportation infrastructure.

Sleep disturbances and chronic psychosocial stress occur at greater rates among individuals experiencing housing instability or economic insecurity than among those who do not experience these conditions. In this review, behavioral risk factors are socially structured and rooted in differences in social, economic, and environmental conditions. Physical activity, diet, sleep, substance use, and stress responses are shaped by neighborhood environments, working conditions, labor organization, food systems, environmental exposures, and access to resources. These influences are distributed inequitably across groups, reflecting long-standing socioeconomic and racial inequities that correspond to consistent differences in cardiovascular risk. When these behaviors are positioned solely as matters of personal responsibility, the structural forces that limit opportunities for cardiovascular health may contribute to disparities that would be obscured. Positioning these factors within a public health and health equity framework, this review incorporates the concept of allostatic load, describing the cumulative physiological "wear and tear" on the body following chronic stress. This concept describes how ongoing external pressures exert cumulative biological effects over time. Repeated stress responses contribute to inflammation, elevated blood pressure, metabolic changes, and disrupted sleep duration, which may elevate cardiovascular risk over time.⁸ By bringing behavioral and environmental risk factors together under an equity lens, these findings may strengthen CVD prevention efforts. Table 1 shows how individual exposures fit into broader structural and systemic settings.

Table 1. Behavioral and Environmental Cardiovascular Risk Domains, Affected Populations, and Policy-Relevant Structural Drivers

Domain of Risk Factors	Primary Exposures or Behaviors for Risks	Most Affected Populations by Risk Factors	Structural Components that Influence Risk Factors	Strategies for Policy and Systems Change
Ambient Air Pollution	Long-term exposure to fine particulate matter (PM _{2.5}), traffic emissions, and industrial pollutants	Low-income communities, communities of color, and residents near highways or industrial zones	Residential segregation, zoning decisions, transportation infrastructure, and weak emissions enforcement	Air quality standards, emissions regulation, land-use planning, and environmental justice enforcement
Contaminated Drinking Water	Arsenic, lead, and agricultural runoff	Rural populations, Indigenous communities, low-income households, and areas with aging infrastructure	Underinvestment in water systems, regulatory gaps, and reliance on groundwater	Water infrastructure investment, regulations
Diet and High-fat, High-Sugar, and High-Salt (HFSS) Foods	High intake of ultra-processed, high-fat, high-sugar, high-salt foods	Low-income households, racial and ethnic minority communities, and food-insecure populations	Food deserts, targeted marketing, pricing, and subsidy structures	Regulation of food marketing, zoning of food outlets, nutrition assistance programs, and pricing policies

Physical Activity	Sedentary behavior, limited access to safe activity spaces	Low-income communities, older adults, women, and people with disabilities	Built environment design, neighborhood safety, transportation access, and work schedules	Walkable urban design, public transit investment, park development, community safety initiatives
Sleep Duration and Insomnia	Short sleep duration, chronic insomnia	Shift workers, individuals with housing instability, and racial and ethnic minority populations	Labor practices, housing conditions, noise exposure, and chronic stress	Work-hour protections, housing stability policies, and neighborhood noise regulation
Psychosocial Stress	Chronic stress exposure, sustained neuroendocrine activation	Communities facing discrimination, economically insecure populations	Racism, economic insecurity, unsafe working conditions, and housing instability	Labor protections, anti-discrimination policy, and community safety investments
Alcohol Consumption	Heavy or harmful alcohol use	Individuals with lower educational attainment, communities exposed to intensive marketing	Alcohol pricing, availability, targeted advertising, and stress exposure	Taxation, advertising restrictions, outlet density regulation, and access to treatment
Tobacco Use	Cigarette smoking, secondhand smoke exposure	Low-income populations, service-sector workers, and communities	Retailer density, marketing practices, and limited smoke-free	Comprehensive smoke-free laws, taxation, retailer density

		with high retailer density	protections	regulation, and cessation access
--	--	----------------------------------	-------------	--

Table 1 (produced by the author) summarizes behavioral and environmental factors that influence cardiovascular risk, showing how individual exposures fit within broader structural and systemic contexts. This framework indicates that cardiovascular health is connected to social environments and highlights policy/system-level changes that may help reduce health inequities.

METHODS

This literature review examines the behavioral and environmental determinants of CVD, with a focus on population trends and health equity. The primary literature search was conducted between January and August 2025. Additional literature searches were conducted during manuscript revisions in 2026 to incorporate new and more relevant evidence. Google Scholar and PubMed were searched using Boolean operators to identify relevant peer-reviewed literature published between 2015 and 2026. The search query combined terms related to cardiovascular outcomes with behavioral and environmental exposures, including: "cardiovascular disease" OR "heart disease" OR "CVD" AND "physical activity" OR "diet" OR "stress" OR "smoking" OR "alcohol" AND "environmental exposures" OR "air pollution" OR "water contamination" OR "neighborhood conditions" OR "food environment". Additional searches linked the terms: "mortality," "incidence," "hypertension," and "coronary artery disease." The selected search terms were chosen to find studies focused on the many modifiable behavioral and ecological factors that are major targets for CVD prevention. The terms "lifestyle" and "CVD prevention" were used to find studies of health-related behaviors, ranging from physical activity, diet, sleep, tobacco use, and alcohol consumption. The term "environmental risk factors" was used to identify studies on structural exposures, such as air pollution, water quality, neighborhood conditions, and food systems. Collectively, these terms align with the theoretical focus of this review by identifying pathways through which repeated exposures and constrained opportunities compound cumulative physiological strain, in relation to the concept of allostatic load.

A subsequent full-text review was performed to determine if studies met the established inclusion criteria. A qualitative synthesis of research evidence was conducted to provide a cohesive narrative, with attention to biological mechanisms and variations in exposure and risk across groups. Studies were included if they were peer-reviewed, examined CVD or cardiovascular risk outcomes, and evaluated at least one behavioral, environmental, or structural exposure within the scope of this review. While studies were excluded if they did not contain cardiovascular-related outcomes, did not address the behavioral or environmental topics evaluated in this review, or were not in the field of CVD prevention.

Titles and abstracts were thoroughly assessed for relevance to the review objectives, along with a full-text analysis of potential studies to determine whether they met the inclusion criteria. Due to the narrative scope of this review, no formal quantitative synthesis of the evidence or standardized risk-of-bias evaluation was conducted. However, several aspects of the study design, including key analytical limitations (i.e., many of the studies in the literature were observational and may have been subject to confounding), were considered when interpreting the results of each study included in this review. Additionally, the peer-reviewed literature is supplemented by information from public health agency reports and international organizations that provide population data and policy and regulatory frameworks, to support the article when evidence was limited in the current body of scholarly research. These sources support contextual and policy background and cannot be used to establish causation between an exposure and an outcome. They are differentiated from empirical research studies throughout the review. Although no formal evidence-grading framework was utilized, the credibility of the findings was evaluated based on study design, sample sizes, methodological limitations, and alignment of findings among the broader literature. The review focuses on key domains (sedentary behavior and physical activity (PA), high-fat, high-sugar, and high-salt (HFSS) and ultra-processed foods, air pollution and CW, sleep duration and insomnia, psychosocial stress and stress hormones, and alcohol and tobacco use) given their potential for modification in CVD prevention and their connection to social and environmental conditions. Hypertension and dyslipidemia (cholesterol) were not treated as primary domains because they function as intermediate clinical risk factors that may occur along with multiple upstream behavioral and structural exposures. Discrimination functions as a major structural stressor that may contribute to varying levels of cardiovascular health.

DISCUSSION

Ambient Air Pollution and Cardiovascular Risk

Air pollution represents a foundational exposure that contributes to cumulative physiological strain through repeated vascular and inflammatory stress.⁹ Ambient air pollution (AAP) is a major risk for CVD, as it is one of the most consequential environmental contributors to cardiovascular morbidity and mortality.¹⁰ Multiple air pollutants contribute to cardiovascular morbidity, particularly particulate matter (PM), a mixture of small airborne particles including dust, dirt, soot, and smoke. Because exposure sources vary, individuals may be exposed to a wide range of pollutants, notably PM_{2.5} (fine particles with a diameter of 2.5 microns or less).¹¹ Exposure to air pollution varies based on where people live, work, and travel. AAP comprises PM and gaseous pollutants (ozone, nitrogen oxides, carbon monoxide, and sulfur dioxide), as well as toxic compounds such as benzene, polycyclic aromatic hydrocarbons, and acrolein.¹² The effect of PM on the heart depends on particle size, measured by the aerodynamic diameter (AD) of the particle.¹⁰ Fine particulate matter (PM_{2.5}) poses cardiovascular risk due to its ability to penetrate the alveolar membranes¹⁰ and enter the body's circulatory system, where it may trigger inflammation and oxidative stress that amplify cardiovascular

mortality. Over time, chronic exposure may culminate in acute cardiovascular events, such as heart attacks and arrhythmias, or premature death.¹³ AAP levels remain persistently elevated worldwide.¹⁴ This determinant may also influence behavioral risk factors, as individuals living near major roadways or industrial areas may be discouraged from outdoor physical activity.¹⁵ In 2019, 99% of the world's population lived in areas that did not meet the World Health Organization (WHO) air quality guidelines.¹⁶ In the United States, almost 79.5 million U.S. residents lived within 500 meters of major roadways in 2020, with pollution concentrations highest within 0-300 meters from the point of release.¹⁷ Globally, around 2.4 billion people are exposed to levels of AAP, with 7 million at risk of premature mortalities each year.¹⁶ Table 2 presents the estimated premature mortality associated with various outdoor air pollution sources in different geographic regions worldwide.

Table 2. Estimated Contributions of Major Outdoor Air Pollution Sources to Premature Mortality Globally and in the United States.

Pollution Source Category	Share of Global Premature Mortality (%)	Share of U.S. Premature Mortality (%)	Regions With Highest Contribution	Dominant Pollutant Constituents
Residential Energy Use	31	6	China, India, Indonesia, Vietnam	Carbon dioxide, carbon monoxide, volatile organic compounds, nitrogen oxides, sulfur dioxide, mercury, fine particulate matter
Agricultural Activities	20	29	Europe, Russia, Japan, eastern United States	Ammonia-derived inorganic fine particulate matter, sulfates, nitrates
Power Generation	14	31	United States, Russia, Korea, Turkey	Sulfate- and nitrate-containing fine particulate matter, mercury

Industrial Sources	7	6	Japan, Germany, China	Sulfate-containing fine particulate matter, volatile organic compounds, and hydrocarbons
Biomass Burning	5	5	Canada, Africa, South America, Australia, Southeast Asia	Fine particulate matter, nitrogen oxides, carbon monoxide, sulfur dioxide, lead, mercury
Land Transportation	5	21	United States, Germany, Russia, Japan	Ultrafine particles, fine particulate matter, nitrogen oxides, volatile organic compounds, ozone
Natural Sources	18	2	Africa, Middle East	Coarse particulate matter, airborne dust

Table 2 illustrates significant regional variations for the primary sources of outdoor air pollution that add to premature deaths in both global and U.S. settings. The estimates presented show the relative contribution of the predominant sources of pollution and demonstrate the geographic variability in how populations are exposed to pollutants. The data were compiled by the author and synthesized from Bhatnagar et al.¹⁷

Evidence linking AAP exposure to cardiovascular morbidity and mortality is substantial, though many of the population-level findings are observational. Thus, there is a possibility that the associations found might be affected by the presence of residual confounding due to such factors as socioeconomic status, occupational and environmental exposure, and preexisting health state. Exposure to AAP is ubiquitous and varies by neighborhood and region. Communities with lower socioeconomic status and communities of color experience disproportionately higher exposure in relation to historical zoning practices, residential segregation, proximity to industrial facilities, and transportation infrastructure.¹⁸ During periods of poor air quality, individuals may lower their exposure by staying indoors, sealing windows, and wearing masks. Ultimately, these findings support population-level and policy-based interventions, including emissions regulations, land-use planning, and enforcement of air quality standards, to advance environmental justice.¹⁹ While individuals may take preventive measures to limit their exposure, lowering cardiovascular risk from air pollution also depends on addressing the conditions that impact exposure throughout multiple communities.

Contaminated Water and Cardiovascular Risk

Chronic exposure to contaminated drinking water could account to long-term physiological strain, increasing cardiovascular vulnerability over the lifespan.²⁰ Arsenic (As) and lead (Pb) are among the most commonly detected contaminants in drinking water today.²¹ Regular consumption of these toxins could be linked to CVD risk and harmful structural heart changes, including left ventricular hypertrophy, defined as thickening of the heart's main pumping chamber (left ventricle).²² These changes impose ongoing stress on the body, heightening susceptibility to CVD. A 2024 study found that over 2 billion people worldwide lack access to clean drinking water,²³ with 1 in 4 people exposed to contaminated water (CW).²⁴ Water contamination may occur when pesticides and fertilizers applied to agricultural lands infiltrate groundwater through rainfall or snowmelt.²⁵ Approximately one million deaths are attributed to unsafe drinking water, lack of sanitation, and poor hygiene.²³ Access to safe drinking water remains unequal across regions, with the greatest burdens falling on low and middle-income countries, particularly in sub-Saharan Africa, where poor infrastructure, limited regulatory oversight, and climate-related stressors increase reliance on contaminated groundwater sources.²⁶ Similar patterns of unequal exposure are seen in the United States. The Strong Heart Study, which evaluated CVD risk factors among American Indian adults, found that they were frequently exposed to arsenic through drinking water in areas where groundwater has been contaminated.²² However, the relationship between contaminated water and cardiovascular risk appears to be different among populations due to the type and concentration of contaminants, exposure duration, and regional water quality. Hence, findings from studies of specific populations, such as the Strong Heart Study, are unlikely to fully represent the cardiovascular effects of contaminated water in other communities. Addressing inequities in exposure to contaminated water requires investment in water infrastructure, regulatory enforcement, and environmental governance, especially in neglected regions, while acknowledging that individual mitigation, such as water quality testing, may also reduce exposure at the household level.²⁷

Physical Activity and Cardiovascular Risk

Constraints on physical activity (PA) operate as upstream conditions that interact with chronic stress and accumulated physiological burden.²⁸ PA is essential for cardiovascular well-being, yet opportunities to be physically active depend on social and environmental conditions. Regular PA offers substantial cardiovascular benefits, as moderate or vigorous activity can strengthen the heart muscle,²⁹ enabling the heart to pump blood more effectively throughout the body.²⁹ The Centers for Disease Control and Prevention (CDC) recommends that adults get 150 minutes of physical activity per week, which is equivalent to 30 minutes a day, five days a week.³⁰ Despite these established guidelines, only 65% of adults worldwide meet recommended activity levels, while 35% fall short.³¹ Meanwhile, increased screen time has exacerbated this trend by encouraging sedentary lifestyles and obesity.³² However, the ability to remain physically active is not merely based on individual effort. It varies based on an individual's environmental conditions, including access to parks

and recreational facilities, work schedules, or income. Several studies have found that parks in low-income neighborhoods are used less than parks in higher-income neighborhoods, even though low-income groups have more leisure time.³³ Consequently, low-income communities, certain racial and ethnic minority groups, women, older adults, and individuals with disabilities are often unable to engage in routine exercise despite similar or greater health needs.³³ Limited access to physical activities may contribute to a sedentary state, defined by prolonged sitting or inactivity,³⁴ doubling the risk of developing or dying from CVD (most prominently CAD), compared to physically active people.³⁵ Because most studies rely on self-reported PA, it remains difficult to determine how PA patterns and underlying environmental and social conditions influence cardiovascular risk. Other factors, including income, pre-existing health conditions, occupation, and neighborhood amenities, could also affect both an individual's ability to engage in PA and their cardiovascular health. Overall, evidence suggests that promoting PA at the population level depends on environments and policies that make active living safer and more attainable, while individual engagement may also minimize CVD risk. Improvements in public transit access, neighborhood safety initiatives, and land-use policies could enable better opportunities for PA at the population level.³⁶

Sleep Duration and Cardiovascular Disease Risk

Insufficient sleep can result from impediments linked to work schedules, housing conditions, and chronic stress, all of which are associated with cardiovascular vulnerability over time.³⁷ Optimal sleep duration is necessary for maintaining cardiovascular health. Sleep specialists recommend that adults get 7-9 hours of sleep each night.³⁸ Transitions to daylight saving time have been linked to short-term changes in myocardial infarction incidence.³⁹ Research has shown that during the first 3-4 days following the springtime shift in March, myocardial infarctions increase by 24% globally.³⁹

Short sleep duration corresponds with multiple adverse health outcomes, including diabetes, hypertension, obesity, and CVD.⁴⁰ Contemporary work schedules, technology use, and daily demands have been linked to delayed sleep timing and insufficient sleep.⁴¹ Sleep is therefore recognized as a social determinant of health, as shift work, racial discrimination, housing instability, and exposure to chronic psychosocial stress may impede opportunities for adequate sleep,⁴² affecting specific communities. In fact, nearly 1 in 3 adults do not meet the recommended seven hours per night.³⁹ Persistent sleep deficiency alters cardiovascular function, straining the cardiovascular system over time.

Insomnia and Cardiovascular Risk

Insomnia is the most prevalent sleep disorder⁴³ and is characterized by difficulty initiating or maintaining sleep.⁴³ Short-term insomnia may occur in response to elevated levels of stress or changes in the environment, typically lasting up to a few days or weeks.⁴⁴ The overall prevalence of insomnia is approximately 22%, with higher rates observed among women than men.⁴⁵ A substantial

body of evidence links insomnia to hypertension, CAD, and HF.⁴⁶ The Nord-Trondelag Health Study followed more than 52,000 adults over 11 years to assess symptoms of insomnia. Depending on the specific insomnia symptom, participants exhibited a 27%-45% higher risk of acute MI.⁴⁶ One study reported that individuals with insomnia had a 45% higher risk of CVD,⁴⁷ and were more likely to have increased stress hormones. Importantly, the relationship between insomnia and CVD is also linked to factors such as age, stress level, lifestyle, and existing health conditions, which means that insomnia does not fully account for a CVD outcome.

Variation in sleep duration and insomnia prevalence is linked to inequities in cardiovascular risk. Black and Hispanic adults in the U.S. who have lower SES levels experience higher rates of short sleep and insomnia; these patterns correlate with their work schedules, housing conditions, and chronic psychosocial stress. The same factors associated with higher rates of cardiovascular morbidity show how social and environmental conditions can influence sleep behaviors.⁴² These trends suggest that policy interventions (i.e., work-hour protections, housing stability measures, and neighborhood noise regulations) may improve cardiovascular well-being, alongside individual-level practices,⁴² such as reducing evening screen time, moderating meal sizes, and maintaining a cool, dark environment. Recommended sleep hours vary among life stages.⁴⁸ Table 3 presents the recommended hours of sleep for individuals at different life stages.

Table 3. Age-Specific Recommended Sleep Duration Across the Life Course

Life Stage	Typical Recommended Daily Duration
Infancy (4–12 months)	Approximately 12–16 hours
Early childhood (1–5 years)	Approximately 10–14 hours
Middle childhood (6–12 years)	Approximately 9–12 hours
Adolescence (13–18 years)	Approximately 8–10 hours
Adulthood (≥ 18 years)	Approximately 7–8 hours

Table 3 outlines the recommended sleep duration among different life stages. The ranges for each age group are based on national recommendations. The data were compiled by the author and synthesized from published national sleep guidelines.⁴⁸

Psychosocial Stress and Neuroendocrine Pathways in Cardiovascular Disease

Psychological stress is an important pathway connecting structural circumstances with cumulative physiological strain, making it a key component of the allostatic load framework.⁴⁹ Psychological

stress activates neuroendocrine pathways that directly alter cardiovascular function. When a threat is perceived, the hypothalamus initiates the body's alert system.⁵⁰ This signaling stimulates the adrenal glands to release stress hormones, particularly adrenaline and cortisol,⁵⁰ which increase heart rate and blood pressure.⁵¹ Excessive release of these hormones can adversely affect the heart. Chronic stress exposure frequently follows broader social and economic inequalities. This is evident in communities facing systemic disadvantage, where economic hardship is associated with intensified stress-related biological wear and tear and greater vulnerability to CVD.⁵²

Adrenaline enhances mental alertness and memory during acute stress.⁵³ However, prolonged exposure has been implicated in endothelial dysfunction and myocardial injury. Excess sympathetic activation is also linked to plaque instability and acute coronary events.⁵⁴ Experimental evidence indicates that adrenaline is involved in plaque rupture, followed by the formation of a thrombus that could completely block blood flow and cause a heart attack.⁵³ Cortisol also influences the cardiovascular system, impacting glucose metabolism, lipid profiles, and blood pressure.⁵⁵ Disruption of sleep and insomnia may amplify these hormonal effects; a recent meta-analysis hypothesized that patients with insomnia have elevated cortisol levels during the day and night, particularly in the evening.⁵⁶

Sustained hormonal activation could exacerbate cardiovascular dysfunction through increased levels of blood sugar, cholesterol, blood pressure, and triglycerides.^{53,57} Chronic activation of hormonal pathways driven by cortisol and adrenaline, is not a universal experience. These problems are tied to environmental inequities: economic insecurity, housing instability, discrimination, and other systemic limitations associated with biological wear and tear. Research has found that chronic psychosocial stressors like racial discrimination are correlated with impaired hypothalamic (pituitary) adrenal axis and altered cortisol secretion.⁵⁸ Due to differences in individual stress responses, chronic stress and CVD risks remains difficult to determine because these outcomes also depend on socioeconomic conditions, lifestyle, and existing health conditions. Moreover, discrimination is a persistent stressor correlated with cumulative biological strain through repeated neuroendocrine activation, a process vital to the concept of allostatic load.⁵⁸ Stress is a pervasive factor that affects nearly all individuals. Although it is experienced at the individual level, many dominant stressors stemming from economic insecurity, unsafe working conditions, and racial discrimination often originate from structural and environmental settings that exceed personal control. Individual-level strategies, particularly regular physical activity, have been associated with lower stress levels and improved cardiovascular health.⁵⁹ Similar benefits are seen with social support networks (family, a close friend, or religious organizations). However, preventive measures alone do not address the structural conditions that generate chronic stress exposure. Successful stress reduction at the population level requires policies and interventions that address structural sources of stress, such as labor protections, community safety initiatives, and housing stability.⁶⁰ Integrating individual stress management with socioenvironmental change is crucial to reducing stress and improving cardiovascular health equity.

Dietary Factors and Cardiovascular Disease Risk

Dietary habits are closely related to food systems and economic access to food, which interact over prolonged periods to create chronic metabolic and inflammatory strain. Diet is a major driver of cardiovascular risk at the population and individual level.⁶¹ Projections estimated that between 2025 and 2050, there will be a 90% increase in CVD prevalence, with an expected 20.5 million CVD deaths in 2025 and 35.6 million CVD deaths in 2050.⁶² Frequent intake of ultra-processed and high-fat, high-sugar, and high-salt (HFSS) foods is tied to premature CVD,⁶³ as these products contain high levels of sodium, saturated and trans fats, and additives such as artificial preservatives, flavor enhancers, and colorants.⁶³ Many low-income neighborhoods have fewer affordable grocery options, more fast-food outlets, and greater marketing of HFSS products.⁶⁴ High densities of fast-food outlets and the targeted advertising of unhealthy products in these areas are associated with greater dietary risks and health inequities, as well as systemic inflammation, insulin resistance, and metabolic syndrome.⁶³ When sustained over time, these dietary exposures act as a chronic physiological stressor, raising cumulative metabolic and inflammatory burden (allostatic load) that increases cardiovascular risk.⁶⁵

High-Fat, High-Sugar, and High-Sodium Dietary Exposures

HFSS foods are products high in saturated and trans fats, added sugars, and sodium, commonly found in processed foods and sweetened beverages.⁶⁶ Overconsumption of HFSS foods is a risk factor for hypertension and deleterious cholesterol levels.⁶⁷ Diets high in trans and saturated fat may raise the risk of heart disease by amplifying low-density lipoprotein (LDL) cholesterol.⁶⁸ Therefore, elevated LDL cholesterol is responsible for atherosclerosis, stroke, CAD, and hypertension.⁶⁹ Sweetened beverages such as sodas, energy drinks, and sports drinks are by far the largest sources of added sugar in the American diet,⁷⁰ with high intake corresponding with insulin resistance and inflammation, thereby increasing cardiometabolic risk.⁷¹ High sodium intake elevates the risk of hypertension, a major risk factor for stroke, heart disease, and impaired renal function.^{72,73} Food insecurity and lack of access to affordable groceries are more prevalent in low-income and disadvantaged communities.⁷⁴ A review of nutrition and health disparities found that among low-income households eligible for SNAP, Black and multiracial households were more likely than White households to experience food insecurity, suggesting structural barriers to accessing benefits.⁷⁴ These disparities highlight that food insecurity is associated to social and environmental conditions, through neighborhood food availability, targeted marketing of dense foods, and socioeconomic constraints, as distinct from individual choice alone. Diet varies widely among individuals, and other factors such as income, culture, eating patterns, and pre-existing health conditions may affect cardiovascular outcome. Although individual dietary choices support cardiovascular health, efforts to reduce cardiovascular risk through nutrition may combine individual-level guidance with population-level strategies (strengthening nutrition assistance programs, regulating unhealthy food marketing, and expanding affordable heart-healthy food options) that improve food environments and decrease structural limitations to healthy eating,⁷⁵ primarily in communities experiencing persistent food insecurity.

Alcohol Use and Cardiovascular Outcomes

Alcohol consumption is widespread and often normalized through cultural, economic, and commercial influences; however, intake exceeding the recommended amount poses serious cardiovascular outcomes.⁷⁶ Excessive alcohol intake is linked to higher blood fats (triglyceride) levels. Increased LDL triglyceride levels are associated with atherosclerotic cardiovascular disease, including myocardial infarction and ischemic stroke.⁷⁷ Other outcomes from overconsumption of alcohol include hypertension, cardiomyopathy, cardiac arrhythmias, or death.⁷⁷ When sustained over years, alcohol exposure interacts with psychosocial and structural stressors in connection with cumulative cardiovascular strain, a characteristic of allostatic load.⁷⁸ Studies report that cardiovascular harm in relation to alcohol use varies with social and environmental conditions spanning from chronic stress, educational attainment, to intensive alcohol marketing. Educational inequities are evident among women with less than a high school education, who experience nearly three times the risk of alcohol-related mortality compared to women holding a bachelor's degree or higher (HR = 2.74). Similar patterns are observed in men, emphasizing that harm from alcohol consumption is concentrated among individuals with lower educational attainment.⁷⁹ Observational studies reported that light to moderate drinkers are at a lower risk for CVD, while heavy, excessive drinkers are at the highest risk.⁸⁰

Since those findings came from observational studies, the apparent lower risk in light/moderate drinkers does not establish that alcohol itself is protective. Therefore, other possible factors could include diet, socioeconomic status, existing health conditions, and other lifestyle behaviors. Public health guidance suggests behavioral strategies such as setting limits, monitoring intake, and addressing triggers.⁸¹ At the same time, population-level strategies, including taxation, advertising restrictions, improved access to care, and harm reduction, are necessary to resolve these inequities.⁸² Alcohol-related cardiovascular harm accumulates alongside biological stress and unequal social conditions; understanding the importance of population-level policies may ease the CVD burden.

Tobacco Use and Cardiovascular Outcomes

Inhaling the toxic mix of chemicals in a cigarette interferes with normal physiological processes. Tobacco toxicants are absorbed into the bloodstream, where they impair cardiovascular function and damage the vascular endothelium.⁸³ Smoking is a leading risk factor for CVD, responsible for over 8 million deaths worldwide annually.⁸⁴ Smoking raises triglycerides, lowers HDL cholesterol, promotes atherosclerotic plaque formation, damages endothelial cells, and is a potential factor in the thickening and narrowing of blood vessels.⁸⁵ Atherosclerotic plaque formations restrict blood flow through arteries and other blood vessels, reducing blood flow to vital organs such as the heart and brain. These processes may result in thrombus formation, with other outcomes such as myocardial infarction, stroke, or death.⁸⁶ Regular exposure to tobacco smoke elevates inflammatory and vascular stress, while cumulative biological strain is associated with cardiovascular vulnerability. The cardiovascular effects of smoking emerge through prolonged biological stress and may be correlated with disparities

in policy protection consistent with an allostatic load framework.⁸⁷ Smoking cessation interventions include behavioral counseling and pharmacologic therapies such as varenicline and bupropion.⁸⁵ These steps may mitigate cardiovascular risk at the individual level, while population-level reductions in tobacco harm could be achieved through tobacco taxation and restrictions on advertising. Tobacco-related cardiovascular harm is disproportionately experienced, with higher burdens concentrated in communities facing economic disadvantages and insufficient access to cessation resources. Workers in low-wage and service-sector occupations are significantly less likely to be covered by comprehensive smokefree laws, primarily in states with local preemption of tobacco control policies. Furthermore, preemption limits local governments' ability to regulate tobacco retailer density, which is often highest in economically disadvantaged regions.⁸⁹ These disparities point to the importance of population-level interventions, through comprehensive smoke-free laws, regulations on retailer density, and equitable access to cessation resources, to decrease exposure and mitigate cardiovascular inequities.

CONCLUSION

Limitations

This review has several methodological limitations. As a narrative literature review, the study was designed to bring together evidence from several behavioral and environmental factors. Although studies were selected based on the inclusion and exclusion criteria described in the Methods, a standardized risk-of-bias assessment was not conducted. Publication bias may also affect the evidence available for this review, since studies finding significant associations are more likely to appear in published literature than studies reporting little or no association. The studies also differ in their populations, locations, methods of measuring exposure, and cardiovascular outcomes. Because of these differences, findings from one study or population may not always apply to another. Another limitation is that much of the evidence discussed in this review comes from observational studies. These studies can identify relationships between exposures and cardiovascular outcomes, but they cannot always determine whether one directly caused the other. Other conditions, including socioeconomic factors, health conditions, and environmental exposures, may also affect these relationships. For these reasons, these constraints may be considered when interpreting the findings.

Implications for Cardiovascular Disease and Health Equity

Cardiovascular disease is a significant threat to global health, responsible for the deaths of millions annually. Although advances have been made in clinical care to enhance treatment options and improve the survival rates for those with CVD, effective strategies to reduce the burden of this disease include both early preventive measures and environmental and public policy-based interventions. Behavioral factors such as diet, exercise, smoking cessation, and stress management play a key role in cardiovascular prevention. However, they are influenced by the contexts and conditions in which people live and work, including housing insecurity, neighborhood pollution, targeted advertising, and structural obstacles to

care, which may influence the distribution of these risk factors across populations. Preventing CVD at a population level requires upstream, equity-driven approaches. Implementation of equitable policies involves ensuring fair access to clean air and water, safe spaces for PA, healthy and affordable food, and culturally competent health services. Public health program initiatives must also prioritize communities that have been marginalized and continue to bear a disproportionate burden of cardiovascular risk. Highlighting the significance of equity-focused interventions and individual strategies is fundamental to reducing cardiovascular disparities and improving population health in a more just and lasting way. Future research studies on this topic would be enhanced by collecting and using exposure data regarding behavioral and environmental exposures with unified frameworks, particularly allostatic load, to quantitatively examine the relationship of cumulative structural conditions and cardiovascular disease risk among diverse populations.

ACKNOWLEDGEMENTS:

I would like to extend my gratitude to the individuals who have supported me in completing this research.

First and foremost, I offer my deepest thanks to my academic mentors and teachers, especially to Debbie Hill, my former 8th-grade English teacher, for their guidance, unwavering support, and the profound insights they provided me throughout this project. Their expertise and feedback were invaluable to my intellectual development.

I am also incredibly grateful to Jacqueline Holstein, my mentor, for her mentorship. Jacqueline's dedicated support and invaluable advice were monumental in shaping this research and helping me navigate the complexities of this topic. Along with Emma Strand, my publishing mentor, for her guidance on navigating the submission process and preparing the manuscript for submission.

Finally, I wish to thank family and friends for their persistent support. They served as a constant source of great strength and motivation, sustaining me throughout this process.

REFERENCES

1. Wang Y, Wang X, Wang C, Zhou J. Global, regional, and national burden of cardiovascular disease, 1990-2021: results from the 2021 global burden of disease study. *Cureus*. 2024 Nov 24;16(11).
2. Jurgens CY, Lee CS, Aycock DM, et al. State of the science: the relevance of symptoms in cardiovascular disease and research: a scientific statement from the american heart association. *Circulation*. 2022;146(12). doi:10.1161/CIR.0000000000001089

3. Di Cesare M, Perel P, Taylor S, et al. The heart of the world. *Glob Heart*. 2024;19(1):11. doi:10.5334/gh.1288
4. Yu G, Gong X, Xu Y, et al. The global burden and trends of four major types of heart disease, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Public Health*. 2023;220:1-9. doi:10.1016/j.puhe.2023.04.005
5. Pampel FC, Krueger PM, Denney JT. Socioeconomic disparities in health behaviors. *Annu Rev Sociol*. 2010;36:349-370. doi:10.1146/annurev.soc.012809.102529
6. Młynarska E, Czarnik W, Fularski P, et al. From atherosclerotic plaque to myocardial infarction—the leading cause of coronary artery occlusion. *IJMS*. 2024;25(13):7295. doi:10.3390/ijms25137295
7. Shilton T, Bauman A, Beger B, et al. More people, more active, more often for heart health – taking action on physical activity. *Global Heart*. 2024;19(1):42. doi:10.5334/gh.1308
8. Logan JG, Barksdale DJ. Allostasis and allostatic load: expanding the discourse on stress and cardiovascular disease. *Journal of Clinical Nursing*. 2008;17(7b):201-208. doi:10.1111/j.1365-2702.2008.02347.x
9. Pryor JT, Cowley LO, Simonds SE. The physiological effects of air pollution: particulate matter, physiology, and disease. *Front Public Health*. 2022;10:882569. Published 2022 July 14. doi:10.3389/fpubh.2022.882569
10. Lee BJ, Kim B, Lee K. Air pollution exposure and cardiovascular disease. *Toxicol Res*. 2014;30(2):71-75. doi:10.5487/TR.2014.30.2.071
11. Oginawati K, Faiqah NSA, Suharyanto, Regia RA, Amin M. Indoor pm_{2.5} and heavy metal composition in blacksmithing factories: a pilot study in Bandung regency, Indonesia. *Urban Science*. 2024;8(4):230. doi:10.3390/urbansci8040230
12. Blaustein JR, Quisel MJ, Hamburg NM, Wittkopp S. Environmental impacts on cardiovascular health and biology: an overview. *Circulation Research*. 2024;134(9):1048-1060. doi:10.1161/circresaha.123.323613
13. U.S. Environmental Protection Agency. Health and Environmental Effects of Particulate Matter (PM). 2025 May 23. <https://www.epa.gov/pm-pollution/health-and-environmental-effects-particulate-matter-pm>
14. World Health Organization (WHO). Billions of People Still Breathe Unhealthy Air: New

- WHO Data. 2022 April 4.
<https://www.who.int/news/item/04-04-2022-billions-of-people-still-breathe-unhealthy-air-ne-w-who-data>
15. U.S. Environmental Protection Agency. Research on Near Roadway and Other Near Source Air Pollution. 2025 February 14.
<https://www.epa.gov/air-research/research-near-roadway-and-other-near-source-air-pollution>
 16. World Health Organization (WHO). Ambient (Outdoor) Air Pollution. 2024 October 24.
[https://www.who.int/news-room/fact-sheets/detail/ambient-\(outdoor\)-air-quality-and-health](https://www.who.int/news-room/fact-sheets/detail/ambient-(outdoor)-air-quality-and-health)
 17. Jaikumar R, Saab GB, Khreis H, et al. Modeling traffic-related air pollution burden of disease using high spatial resolution data. *Journal of Transport and Health*. 2025;44:102155.
doi:10.1016/j.jth.2025.102155
 18. Woo B, Kravitz-Wirtz N, Sass V, Crowder K, Teixeira S, Takeuchi DT. Residential segregation and racial/ethnic disparities in ambient air pollution. *Race Soc Probl*. 2019;11(1):60-67. doi:10.1007/s12552-018-9254-0
 19. Giles LV, Barn P, Künzli N, et al. From good intentions to proven interventions: effectiveness of actions to reduce the health impacts of air pollution. *Environ Health Perspect*. 2011;119(1):29-36. doi:10.1289/ehp.1002246
 20. Evans E, Jacobs M, Fuller D, Hegland K, Ellis C. Allostatic load and cardiovascular disease: a systematic review. *Am J Prev Med*. 2025;68(6):1072-1079.
doi:10.1016/j.amepre.2025.02.016
 21. Jennrich P. The influence of arsenic, lead, and mercury on the development of cardiovascular diseases. *ISRN Hypertension*. 2013;2013:1-15. doi:10.5402/2013/234034
 22. Pichler G, Grau-Perez M, Tellez-Plaza M, et al. Association of arsenic exposure with cardiac geometry and left ventricular function in young adults: evidence from the strong heart family study. *Circ: Cardiovascular Imaging*. 2019;12(5):e009018.
doi:10.1161/CIRCIMAGING.119.009018
 23. World Health Organization (WHO). Drinking Water. 2023 September 13.
<https://www.who.int/news-room/fact-sheets/detail/drinking-water>
 24. Ritchie H, Spooner F, Roser M. Clean water: Our World in Data. 2019 Sep 26.
<https://ourworldindata.org/clean-water>

25. U.S. Geological Survey. Pesticides in Groundwater. 2018 October 9.
<https://www.usgs.gov/special-topics/water-science-school/science/pesticides-groundwater>
26. Greenwood EE, Lauber T, van den Hoogen J, et al. Mapping safe drinking water use in low- and middle-income countries. *Science*. 2024;385(6710):784-790.
doi:10.1126/science.adh9578
27. VanDerslice J. Drinking water infrastructure and environmental disparities: evidence and methodological considerations. *American Journal of Public Health*. 2011;101(S1):S109-S114.
doi:10.2105/ajph.2011.300189
28. Robertson T, Benzeval M, Whitley E, Popham F. The role of material, psychosocial, and behavioral factors in mediating the association between socioeconomic position and allostatic load (measured by cardiovascular, metabolic, and inflammatory markers). *Brain Behavior and Immunity*. 2014;45:41-49. doi:10.1016/j.bbi.2014.10.005
29. Dhuli K, Naureen Z, Medori MC, et al. Physical activity for health. *J Prev Med Hyg*. 2022;63(2 Suppl 3):E150-E159. doi:10.15167/2421-4248/jpmh2022.63.2S3.2756
30. Piercy KL, Troiano RP, Ballard RM, et al. The physical activity guidelines for americans. *JAMA*. 2018;320(19):2020. doi:10.1001/jama.2018.14854
31. World Health Organization: WHO. Nearly 1.8 billion adults at risk of disease from not doing enough physical activity. World Health Organization. 2024 June 26.
<https://www.who.int/news/item/26-06-2024-nearly-1.8-billion-adults-at-risk-of-disease-from-not-doing-enough-physical-activity>
<https://www.who.int/news/item/26-06-2024-nearly-1.8-billion-adults-at-risk-of-disease-from-not-doing-enough-physical-activity>
32. Ramsey Buchanan L, Rooks-Peck CR, Finnie RKC, et al. Reducing recreational sedentary screen time: a community guide systematic review. *Am J Prev Med*. 2016;50(3):402-415.
doi:10.1016/j.amepre.2015.09.030
33. Cohen DA, Han B, Park S, Williamson S, Derosé KP. Park Use and Park-Based Physical Activity in Low-Income Neighborhoods. *J Aging Phys Act*. 2019;27(3):334-342.
doi:10.1123/japa.2018-0032
34. DiPietro L, Al-Ansari SS, Biddle SJH, et al. Advancing the global physical activity agenda: recommendations for future research by the 2020 WHO physical activity and sedentary behavior guidelines development group. *Int J Behav Nutr Phys Act*. 2020;17(1):143.
doi:10.1186/s12966-020-01042-2

35. Press V, Freestone I, George CF. Physical activity: the evidence of benefit in the prevention of coronary heart disease. *QJM*. 2003;96(4):245-251. doi:10.1093/qjmed/hcg041
36. Zhu X, Ory MG, Xu M, et al. Physical activity impacts of an activity-friendly community: a natural experiment study protocol. *Front Public Health*. 2022;10:929331. Published 2022 June 15. doi:10.3389/fpubh.2022.929331
37. Pan Y, Zhou Y, Shi X, He S, Lai W. The association between sleep deprivation and the risk of cardiovascular diseases: A systematic meta-analysis. *Biomed Rep*. 2023;19(5):78. Published 2023 September 12. doi:10.3892/br.2023.1660
38. St-Onge M, Jelic S. Maintaining stable, adequate sleep: instrumental to cardiometabolic health. *JAHA*. 2025;14(15):e043464. doi:10.1161/JAHA.125.043464
39. Li J, Cao D, Huang Y, et al. Sleep duration and health outcomes: an umbrella review. *Sleep Breath*. 2022;26(3):1479-1501. doi:10.1007/s11325-021-02458-1
40. St-Onge MP, Aggarwal B, Fernandez-Mendoza J, Johnson D, Kline CE, Knutson KL, Redeker N, Grandner MA, American Heart Association Council on Lifestyle and Cardiometabolic Health, Council on Cardiovascular and Stroke Nursing, Council on Clinical Cardiology, and Council on Quality of Care and Outcomes Research. Multidimensional sleep health: definitions and implications for cardiometabolic health: A scientific statement from the American Heart Association. *Circulation: Cardiovascular Quality and Outcomes*. 2025 May;18(5):e000139.
41. White KM, Barber LE, Shipman-Lacewell JR, Azeez O, Collop N, Johnson DA. Understanding and addressing social determinants to advance sleep health equity in the united states: a blueprint for research, practice, and policy. *Curr Sleep Medicine Rep*. 2025;11(1):30. doi:10.1007/s40675-025-00339-7
42. Jackson CL, Redline S, Emmons KM. Sleep as a potential fundamental contributor to disparities in cardiovascular health. *Annu Rev Public Health*. 2015;36:417-440. doi:10.1146/annurev-publhealth-031914-122838
43. McCarthy CE. Sleep disturbance, sleep disorders and co-morbidities in the care of the older person. *Medical Sciences*. 2021;9(2):31. doi:10.3390/medsci9020031
44. Krystal AD, Ashbrook LH, Prather AA. What is insomnia? *JAMA*. 2021;326(23):2444. doi:10.1001/jama.2021.19283
45. Zeng LN, Zong QQ, Yang Y, et al. Gender difference in the prevalence of insomnia: a meta-analysis of observational studies. *Front Psychiatry*. 2020;11:577429.

doi:10.3389/fpsy.2020.577429

46. Javaheri S, Redline S. Insomnia and Risk of Cardiovascular Disease. *Chest*. 2017;152(2):435-444. doi:10.1016/j.chest.2017.01.026
47. Andersen ML, Poyares D, Tufik S. Insomnia and cardiovascular outcomes. *Sleep Sci*. 2021;14(1):1-2. doi:10.5935/1984-0063.20200109
48. Mireku MO, Rodriguez A. Sleep duration and waking activities in relation to the national sleep foundation's recommendations: an analysis of us population sleep patterns from 2015 to 2017. *IJERPH*. 2021;18(11):6154. doi:10.3390/ijerph18116154
49. Palix C, Chauveau L, Felisatti F, et al. Allostatic load, a measure of cumulative physiological stress, impairs brain structure but not β -accumulation in older adults: an exploratory study. *Frontiers in Aging Neuroscience*. 2025;17:1508677. doi:10.3389/fnagi.2025.1508677
50. Mayo Clinic. Chronic stress puts your health at risk. *Stress Management*. 2023 August 1. <https://www.mayoclinic.org/healthy-lifestyle/stress-management/in-depth/stress/art-20046037>
51. Chu B, Marwaha K, Sanvictores T, Awosika AO, Ayers D. Physiology, stress reaction. *InStatPearls*. StatPearls Publishing. 2024 May 7. <https://www.ncbi.nlm.nih.gov/books/NBK541120/>
52. Schultz WM, Kelli HM, Lisko JC, et al. Socioeconomic Status and Cardiovascular Outcomes: Challenges and Interventions. *Circulation*. 2018;137(20):2166-2178. doi:10.1161/CIRCULATIONAHA.117.029652
53. Schwabe L. Memory under stress: from adaptation to disorder. *Biological Psychiatry*. 2025;97(4):339-348. doi:10.1016/j.biopsych.2024.06.005
54. Rezkalla SH, Kloner RA. Stress and acute coronary syndrome. *Cardiol Res*. 2025;16(6):467-474. doi:10.14740/cr2123
55. The effects of stress on the heart. <https://www.medstarhealth.org/blog/heart-and-stress>
56. Dressle RJ, Feige B, Spiegelhalder K, et al. HPA axis activity in patients with chronic insomnia: A systematic review and meta-analysis of case-control studies. *Sleep Med Rev*. 2022;62(101588):101588. doi:10.1016/j.smrv.2022.101588
57. Tsai SY, Hsu JY, Lin CH, et al. Association of stress hormones and the risk of cardiovascular

- diseases systematic review and meta-analysis. *International Journal of Cardiology Cardiovascular Risk and Prevention*. 2024;23:200305. doi:10.1016/j.ijcrp.2024.200305
58. Lee DB, Peckins MK, Heinze JE, Miller AL, Assari S, Zimmerman MA. Psychological pathways from racial discrimination to cortisol in African American males and females. *J Behav Med*. 2018;41(2):208-220. doi:10.1007/s10865-017-9887-2
59. University of Rochester Medical Center. Stress can increase your risk for heart disease. *Health Encyclopedia*. 2025.
<https://www.urmc.rochester.edu/encyclopedia/content?ContentTypeID=1&ContentID=2171>
60. Kirkbride JB, Anglin DM, Colman I, et al. The social determinants of mental health and disorder: evidence, prevention and recommendations. *World Psychiatry*. 2024;23(1):58-90. doi:10.1002/wps.21160
61. Wang W, Liu Y, Li Y, et al. Dietary patterns and cardiometabolic health: Clinical evidence and mechanism. *MedComm*. 2023;4(1):e212. doi:10.1002/mco2.212
62. Chong B, Jayabaskaran J, Jauhari SM, Chan SP, Goh R, Kueh MT, Li H, Chin YH, Kong G, Anand VV, Wang JW. Global burden of cardiovascular diseases:projections from 2025 to 2050. *European Journal of Preventive Cardiology*. 2025 Aug;32(11):1001-15. doi:10.1093/eurjpc/zwae281
63. Willett Y, Yang C, Dunn J, et al. Consumption of ultra-processed foods and increased risks of cardiovascular disease in u. S. Adults. *The American Journal of Medicine*. Published online January 2026:S0002934326000549. doi:10.1016/j.amjmed.2026.01.012
64. Hernández-Caravaca I, Martínez-García A, Trescastro-López EM, Plaza-Gavaldón Á, Martí-Cremades J, Moncho J. Influence of socioeconomic status on the retailfood environment in alicante. *Nutrients*. 2024;16(23):4127. doi:10.3390/nu16234127
65. Babalola OO, Akinnusi E, Ottu PO, et al. The impact of ultra-processed foods on cardiovascular diseases and cancer: Epidemiological and mechanistic insights.Aspects of *Molecular Medicine*. 2025;5:100072. doi:10.1016/j.amolm.2025.100072
66. Vadiveloo MK, Gardner CD, Bleich SN, et al. Ultraprocessed foods and their association with cardiometabolic health: evidence, gaps, and opportunities: a science advisory from the american heart association. *Circulation*. 2025;152(12). doi:10.1161/CIR.0000000000001365
67. Thapsuwan S, Phulkerd S, Chamrathirong A, et al. Relationship between consumption of high fat, sugar or sodium (Hfss) food and obesity and non-communicable diseases. *BMJNPH*.

- 2024;7(1):78-87. doi:10.1136/bmjnph-2023-000794Maki KC, Dicklin MR,
68. Kirkpatrick CF. Saturated fats and cardiovascular health: Current evidence and controversies. *Journal of Clinical Lipidology*. 2021;15(6):765-772. doi:10.1016/j.jacl.2021.09.049
69. Mortensen MB, Dzaye O, Bøtker HE, et al. Low-density lipoprotein cholesterol is predominantly associated with atherosclerotic cardiovascular disease events in patients with evidence of coronary atherosclerosis: the western denmark heart registry. *Circulation*. 2023;147(14):1053-1063. doi:10.1161/CIRCULATIONAHA.122.061010
70. Ooi JY, Wolfenden L, Sutherland R, et al. A systematic review of the recent consumption levels of sugar-sweetened beverages in children and adolescents from the world health organization regions with high dietary-related burden of disease. *Asia Pac J Public Health*. 2022;34(1):11-24. doi:10.1177/10105395211014642
71. Santos LP, Gigante DP, Delpino FM, Maciel AP, Bielemann RM. Sugar sweetened beverages intake and risk of obesity and cardiometabolic diseases in longitudinal studies: A systematic review and meta-analysis with 1.5 million individuals. *Clinical Nutrition ESPEN*. 2022;51:128-142. doi:10.1016/j.clnesp.2022.08.021
72. Surma S, Francuz T, Ciechanowski K. High salt intake and cardiovascular disease: an overview. In: Preedy VR, Patel VB, eds. *Handbook of Public Health Nutrition*. Springer Nature Switzerland; 2025:1-25. doi:10.1007/978-3-031-32047-7_190-1
73. Cappuccio FP. Cardiovascular and other effects of salt consumption. *Kidney Int Suppl* (2011). 2013;3(4):312-315. doi:10.1038/kisup.2013.65
74. Samuel LJ, Crews DC, Swenor BK, et al. Supplemental Nutrition Assistance Program access and racial disparities in food insecurity. *JAMA Netw Open*. 2023;6(6):e2320196. doi:10.1001/jamanetworkopen.2023.20196
75. Anand SS, Hawkes C, de Souza RJ, et al. Food consumption and its impact on cardiovascular disease: importance of solutions focused on the globalized food system.a report from the Workshop Convened by the World Heart Federation. *J Am Coll Cardiol*. 2015;66(14):1590-1614. doi:10.1016/j.jacc.2015.07.050
76. British Heart Foundation. Effects of alcohol on your heart. British Heart Foundation. 2025 June 3.
<https://www.bhf.org.uk/information-support/heart-matters-magazine/medical/effects-of-alcohol-on-your-heart>.

77. Balling M, Afzal S, Davey Smith G, et al. Elevated ldl triglycerides and atherosclerotic risk. *Journal of the American College of Cardiology*. 2023;81(2):136-152. doi:10.1016/j.jacc.2022.10.019
78. Rogers JM, Epstein DH, Phillips K, Strickland JC, Preston KL. Exploring the relationship between substance use and allostatic load in a treatment/research cohort and in a U.S. Probability Sample (NHANES 2009-2016). *Front Psychiatry*. 2021;12:630195. Published 2021 August 2. doi:10.3389/fpsy.2021.630195
79. Andreacchi AT, Hobin E, Siddiqi A, Smith BT. Socioeconomic inequities in alcohol-attributable mortality by sex/gender and age in Canada: a 13-year population-representative cohort study. *American Journal of Epidemiology*. 2024;194(11):3069-3078. doi:10.1093/aje/kwae385
80. Lavie CJ, O'Keefe JH, Bhatti SK, Bajwa A, DiNicolantonio JJ. Alcohol and cardiovascular health: The dose makes the poison ... or the remedy. *Mayo Clinic*. 2014;89(3):382-393. doi:10.1016/j.mayocp.2013.11.005
81. Centers for Disease Control and Prevention. Getting Started with Drinking Less. Drink Less, Be Your Best. 2024 October 8. <https://www.cdc.gov/drink-less-be-your-best/getting-started-with-drinking-less/index.html>
82. Gilmore W, Gilmore I. Preventing alcohol-related harm: effective strategies and the role of health professionals. *Br J Hosp Med (Lond)*. 2025;86(8):1-12. doi:10.12968/hmed.2024.0719
83. Münzel T, Crea F, Rajagopalan S, Lüscher T. Nicotine and the cardiovascular system: unmasking a global public health threat. *European Heart Journal*. Published online December 18, 2025:ehaf1010. doi:10.1093/eurheartj/ehaf1010
84. Cho JH, Shin SY, Kim H, et al. Smoking cessation and incident cardiovascular disease. *JAMA Network Open*. 2024;7(11):e2442639-e2442639. doi:https://doi.org/10.1001/jamanetworkopen.2024.42639
85. Klein LW. Pathophysiologic mechanisms of tobacco smoke producing atherosclerosis. *Current Cardiology Reviews*. 2022;18(6):60-67. doi:10.2174/1573403X18666220411113112
86. He Z, Luo J, Lv M, et al. Characteristics and evaluation of atherosclerotic plaques: an overview of state-of-the-art techniques. *Front Neurol*. 2023;14. doi:10.3389/fneur.2023.1159288
87. Rahman M, Alatiqi M, Al Jarallah M, et al. Cardiovascular effects of smoking and smoking

- cessation: a 2024 update. *Glob Heart*. 2025;20(1):15. 2025 February 19. doi:10.5334/gh.1399
88. Centers for Disease Control and Prevention. Health Effects of Cigarettes: Cardiovascular Disease. 2024 September 17. Available at:
<https://www.cdc.gov/tobacco/about/cigarettes-and-cardiovascular-disease.html>
89. Office of the Surgeon General. U.S. Department of Health and Human Services. Promising interventions to reduce tobacco-related health disparities. In *Eliminating Tobacco-Related Disease and Death: Addressing Disparities: A Report of the Surgeon General*. 2024.
<https://www.ncbi.nlm.nih.gov/books/NBK614483/>